

# Role of H<sub>2</sub>O<sub>2</sub> in pea seed germination

Gregorio Barba-Espín, José Antonio Hernández\* and Pedro Diaz-Vivancos

Grupo de Biotecnología de Frutales; CEBAS-CSIC; Campus Espinardo; Murcia, Spain

**Keywords:** hydrogen peroxide, redox signaling, germination, seed priming, dormancy

The imbibition of pea seeds with hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) increased the germination as well as the seedling growth, producing an invigoration of the seeds. We propose that H<sub>2</sub>O<sub>2</sub> could act as signaling molecule in the beginning of seed germination involving specific changes at proteomic, transcriptomic and hormonal levels. These findings have practical implication in the context of seed priming technologies to invigorate low vigour seeds.

Germination process is associated with many metabolic, cellular, and molecular events, coordinated by a complex regulatory network. The reactive oxygen species (ROS) production by germinating seeds has often been considered as a negative effect that might affect the germination process, but provided that their accumulation is tightly regulated by the balance between production and scavenging, these toxic molecules now appear as being beneficial for germination.<sup>1</sup>

Strategies for improving the growth and development of crop species have been investigated for many years. Seed priming is a pre-sowing strategy to influence seed germination and seedling development by modulating pre-germination metabolic activity prior to emergence of the radicle and generally enhances germination rate and plant performance.<sup>2,3</sup> From a biochemical and molecular point of view, studying germination is difficult because a population of seeds does not complete the process synchronously.<sup>4</sup> Seed priming has been found as technology to enhance rapid and uniform emergence, and to achieve high vigour and better yields. This process generally causes faster germination and faster field emergence, which has practical agronomic implications, notably under adverse germination conditions.<sup>5</sup>

## Effect of ROS-Related Compound in Seed Germination and Seedling Growth

We have tested the priming effect of several compounds on pea seeds. The assayed compound can be divided in three major groups: directly involved in the antioxidative metabolism [hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), reduced and oxidized glutathione (GSH, GSSG) and ascorbic acid (ASC)]; compounds related with the amino acid cysteine [N-acetylcysteine (NAC), thioproline (TP) and L-2-oxo-4-thiazolidine-carboxylic acid (OTC)]; and compounds that trigger the systemic acquired resistance in plants [salicylic acid (SA) and its functional analog benzothiadiazole (BTH)].


The tripeptide thiol glutathione (GSH) is at the hub of the complex antioxidant networks of plant and animal cells, where it

participates in the cellular redox signaling networks that influence growth, development and defense.<sup>6,7</sup> It has long been recognized that ASC also exert a strong influence on plant growth and development.<sup>8</sup> The GSH synthesis is regulated by cysteine availability, thus compounds increasing its contents could produce an increase of GSH levels. In this sense, OTC is an artificial cysteine precursor whereas TP is a proline analog that is converted by proline oxidase to N-formyl-L-cysteine which is presumably hydrolyzed to cysteine.<sup>9</sup>

To carry out this work, pea seeds were imbibed for 24 h in dH<sub>2</sub>O or in the compounds described above at different concentrations. Seeds were then washed twice with dH<sub>2</sub>O and placed in Petri dishes with two layers of filter paper moistened with dH<sub>2</sub>O. Seeds were incubated at 25°C for 48 h in darkness, in a Cooled Incubator (MIR-153 Sanyo).

Figure 1 shows that, except H<sub>2</sub>O<sub>2</sub>, none of the assayed compounds had a positive effect on seed germination or seedling growth (measured as fresh weight and length). Exogenous H<sub>2</sub>O<sub>2</sub> showed a priming effect in the germination of pea seeds in a concentration dependent-manner obtaining more vigorous seedlings, being 20 mM H<sub>2</sub>O<sub>2</sub> the concentration that produced the best response in terms of growth (Fig. 1). The increase in seedling growth by 20 mM H<sub>2</sub>O<sub>2</sub> was also evident 24 h after imbibition (Fig. 2). The priming effect of H<sub>2</sub>O<sub>2</sub> was also noticeable at shorter times of imbibition (Table 1). After 12 h of imbibition about 15% of seeds had germinated; however, no germination occurred at this time in seeds imbibed in water. At 24 h of imbibition, this percentage had reached nearly 75%, whereas control seed germination remained at low level (Table 1). In preliminary experiments, we noticed that H<sub>2</sub>O<sub>2</sub> concentrations higher than 20 mM (40 to 100 mM) also stimulated the germination rate after 24 h imbibition. However, at short-term of post-imbibition (6 h) we observed that these H<sub>2</sub>O<sub>2</sub> levels induced a pronounced curvature as well as an abnormal growth of the radicle at 24 h and 48 h post-imbibition (Fig. 2). In addition, H<sub>2</sub>O<sub>2</sub> concentrations above 100 mM reduced the pea seeds germination rate (data not shown).

\*Correspondence to: José Antonio Hernández; Email: jahernan@cebas.csic.es  
Submitted: 10/27/11; Revised: 11/28/11; Accepted: 11/28/11  
<http://dx.doi.org/10.4161/psb.18881>




**Figure 1.** Effect of imbibition with different compounds on the percentage of germination of pea seeds (A) and on the fresh weight (B) and length (C) of the seedlings 48 h after imbibition. Used treatments concentrations (in mM): H<sub>2</sub>O<sub>2</sub> (5, 10 and 20); GSH (0.125, 0.250 and 0.500); GSSG (0.0625, 0.125, 0.250); ASC, OTC, NAC, SA and BTH (0.25, 0.50 and 1). Black, gray and dark gray bars correspond to the lower, intermediate and the higher concentrations of each treatment, respectively. Different letters indicate statistical significance according to Tukey's test ( $p < 0.05$ ).

Interestingly, we observed a differential response depending on when H<sub>2</sub>O<sub>2</sub> was supplied. When 5 mM H<sub>2</sub>O<sub>2</sub> was added during the incubation in plates, after imbibition in dH<sub>2</sub>O, the percentage germination was similar to that of control seeds. In contrast, the presence of 10 or 20 mM H<sub>2</sub>O<sub>2</sub> in Petri dishes produced a negative effect on germination (Fig. 3).

### H<sub>2</sub>O<sub>2</sub> Signaling during Seed Germination

Numerous recent works show now that ROS would play a key signaling role in the achievement of major events of seed life, such as germination or dormancy release. In fact, hydrogen peroxide, nitric oxide, hydroxyl radicals and superoxide radicals have been shown to accumulate during seed germination in various species.<sup>1</sup> Many works have reported that exogenous application of H<sub>2</sub>O<sub>2</sub>




**Figure 2.** Effect of H<sub>2</sub>O<sub>2</sub> imbibition (0, 20, 40 and 80 mM) on early pea seedling growth after incubation in darkness.

can improve seed germination in many plant species.<sup>10-12</sup> The interplay between ROS and hormone signaling pathways lead to changes in gene expression or in cellular redox status that would play a role in the perception of environmental factors by seeds during their germination.<sup>1</sup> Recently we have shown that H<sub>2</sub>O<sub>2</sub> coordinates the beginning of pea seed germination, acting as a priming factor that involves specific changes at proteome, transcriptome and hormonal levels, resulting in an acceleration of the germination process most probably due to invigoration of the seeds.<sup>13</sup> H<sub>2</sub>O<sub>2</sub> would induce a MAPK-dependent decrease in abscisic acid (ABA) contents in the seed as well as the carbonylation of seed storage proteins,<sup>13</sup> favoring their mobilization, and some glycolytic enzymes that could stimulate the phosphate pentose pathway (oxPPP).<sup>14</sup> The oxPPP activation could provide NADPH for the thioredoxin system, involved in seed germination

**Table 1.** Effect of 20 mM H<sub>2</sub>O<sub>2</sub> imbibition on the percentage of pea seed germination at different time periods. Four batches of 25 seeds for each treatment were used to calculate the rate of germination

	Time (h)	Water	H <sub>2</sub> O <sub>2</sub>
Imbibition	6 h	0	0
	12 h	0	14
	24 h	15	75



**Figure 3.** Percentage of germination of pea seeds after imbibition in the presence of dH<sub>2</sub>O (0 h) and 48 h of incubation in plates in the presence of different H<sub>2</sub>O<sub>2</sub> concentrations.

and seedling development.<sup>15</sup> Alternatively, H<sub>2</sub>O<sub>2</sub> could act, directly or indirectly impairing the ABA transport from the cotyledon to the embryo inducing a decrease in ABA, stimulating the germination process.<sup>13</sup> Finally, the decrease in ABA could induce a MAPK-mediated decrease in the ethylene precursor 1-aminocyclopropane carboxylic acid, favoring epicotyl and radicle emergence by H<sub>2</sub>O<sub>2</sub> treatment.<sup>13</sup>

## References

- Bailly C, El-Maarouf-Bouteau H, Corbineau F. From intracellular signaling networks to cell death: the dual role of reactive oxygen species in seed physiology. *C R Biol* 2008; 331:806-14; PMID:18926495; <http://dx.doi.org/10.1016/j.crv.2008.07.022>
- Bradford KJ. Manipulation of seed water relations via osmotic priming to improve germination under stress conditions. *Hort Sci* 1986; 21:1105-12.
- Taylor AG, Harman GE. Concepts and technologies of selected seed treatments. *Annu Rev Phytopathol* 1990; 28:321-39; <http://dx.doi.org/10.1146/annurev.py.28.090190.001541>
- Still DW, Dahal P, Bradford KJ. A single-seed assay for endo- $\beta$ -mannanase activity from tomato endosperm and radicle tissues. *Plant Physiol* 1997; 113:13-20; PMID:12223589
- Black M, Bewley JD. *Seed Technology and Its Biological Basis*. Sheffield Academic Press Ltd., Sheffield 2000.
- Foyer CH, Noctor G. Redox homeostasis and antioxidant signaling: a metabolic interface between stress perception and physiological responses. *Plant Cell* 2005; 17:1866-75; PMID:15987996; <http://dx.doi.org/10.1105/tpc.105.033589>
- Maughan S, Foyer CH. Engineering and genetic approaches to modulating the glutathione network in plants. *Physiol Plant* 2006; 126:382-97; <http://dx.doi.org/10.1111/j.1399-3054.2006.00684.x>

- Pignocchi C, Foyer CH. Apoplastic ascorbate metabolism and its role in the regulation of cell signalling. *Curr Opin Plant Biol* 2003; 6:379-89; PMID:12873534; [http://dx.doi.org/10.1016/S1369-5266\(03\)00069-4](http://dx.doi.org/10.1016/S1369-5266(03)00069-4)
- Hilton JL, Pillai P, Norman HA. Fate of safening agent L-2-oxothiazolidine-4-carboxylic acid in shorgum (*Sorghum bicolor*). *Weed Sci* 1990; 38:201-5.
- Fontaine O, Huault C, Pavis N, Billard JP. Dormancy breakage of *Hordeum vulgare* seeds: Effect of hydrogen peroxide and stratification on glutathione level and glutathione reductase activity. *Plant Physiol Biochem* 1994; 32:677-83.
- Katzman LS, Taylor AG, Langhans RW. Seed enhancements to improve spinach germination. *Hort Sci* 2001; 36:979-81.
- Zeinalabedini M, Majourhat K, Hernández JA, Martínez-Gómez P. Breaking seed dormancy in long-term stored seeds from Iranian wild almond species. *Seed Sci Technol* 2009; 37:267-75.
- Barba-Espín G, Diaz-Vivancos P, Job D, Belghazi M, Job C, Hernández JA. Understanding the role of H<sub>2</sub>O<sub>2</sub> during pea seed germination: a combined proteomic and hormone profiling approach. *Plant Cell Environ* 2011; 34:1907-19; PMID:21711356; <http://dx.doi.org/10.1111/j.1365-3040.2011.02386.x>

- Job C, Rajjou L, Lovigny Y, Belghazi M, Job D. Patterns of protein oxidation in Arabidopsis seeds and during germination. *Plant Physiol* 2005; 138:790-802; PMID:15908592; <http://dx.doi.org/10.1104/pp.105.062778>
- Lozano RM, Wong JH, Yee BC, Peters A, Kobrehel K, Buchanan BB. New evidence for a role for thioredoxin h in germination and seed development. *Planta* 1996; 200:100-6; <http://dx.doi.org/10.1007/BF00196655>
- Barba-Espín G, Diaz-Vivancos P, Clemente-Moreno MJ, Albacete A, Faize L, Faize M, et al. Interaction between hydrogen peroxide and plant hormones during germination and the early growth of pea seedlings. *Plant Cell Environ* 2010; 33:981-94; PMID:20102539; <http://dx.doi.org/10.1111/j.1365-3040.2010.02120.x>
- Harris D, Joshi A, Khan PA, Gothkar P, Sodhi PS. On-farm seed priming in semi-arid agriculture: Development and evaluation in maize, rice and chickpea in India using participatory methods. *Exp Agric* 1999; 35:15-29; <http://dx.doi.org/10.1017/S0014479799001027>

As mentioned above, the H<sub>2</sub>O<sub>2</sub> treatment also stimulates the early growth of pea seedlings (Table 1). Previously, we have described that the H<sub>2</sub>O<sub>2</sub>-induced increase in pea seedling growth was correlated with the induction of proteins related to plant growth, cellular signaling and cell cycle control (14–3-3 protein, profilin, proteasome, translationally-controlled tumor protein), as well as with a substantial decrease in the levels of the hormones ABA and zeatin-riboside (ZR).<sup>16</sup> Moreover, a decrease of a polypeptide with homology to an ABA-responsive protein was observed, suggesting an interplay between the redox state and plant hormones, orchestrated by H<sub>2</sub>O<sub>2</sub>, in the induction of proteins related to plant signaling and development during the early growth of pea seedlings.<sup>16</sup> The molecular mechanisms implicated in this differential response remain unclear. They are of considerable interest, notably in the context of improving crop yields by invigoration seed treatments both in commercial applications<sup>2</sup> and in developing countries.<sup>17</sup>

## Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

## Acknowledgments

This work was supported by Fundación Séneca—Agencia de Ciencia y Tecnología de la Región de Murcia (projects 05571/PI/07 and 11883/PI/09). G.B.-E. thanks CSIC for his JAE research fellowship. P.D.-V. thanks CSIC for his JAEdoc research contract.